

## **SCAI/AATS/ACCF/STS Multisociety Expert Consensus Statement:**

### **Operator & Institutional Requirements for**

### **Transcatheter Valve Repair and Replacement; Part 1 TAVR**

Carl L. Tommaso, MD<sup>1</sup>, R. Morton Bolman III<sup>2</sup>, MD, Ted Feldman, MD<sup>3</sup>, Joseph Bavaria, MD<sup>4</sup>, Michael A. Acker, MD<sup>4</sup>, Gabriel Aldea, MD<sup>5</sup>, Duke E. Cameron, MD<sup>6</sup>, Larry S. Dean, MD<sup>5</sup>, David Fullerton, MD<sup>7</sup>, Ziyad M. Hijazi, MD<sup>8</sup>, Eric Horlick, MD<sup>9</sup>, D. Craig Miller, MD<sup>10</sup>, Marc R. Moon, MD<sup>11</sup>, Richard Ringel, MD<sup>6</sup>, Carlos E. Ruiz, MD, PhD<sup>12</sup>, Alfredo Trento, MD<sup>13</sup>, Bonnie H. Weiner MD<sup>14</sup>, Evan M. Zahn, MD<sup>15</sup>

<sup>1</sup>Skokie Hospital NSUHS, <sup>2</sup> Harvard Medical School, <sup>3</sup> Evanston Hospital, NSUHS, <sup>4</sup>University of Pennsylvania Medical Center, <sup>5</sup>University of Washington Medical Center, <sup>6</sup>The Johns Hopkins Hospital, <sup>7</sup>University of Colorado, <sup>8</sup>Rush University Medical Center, <sup>9</sup>Toronto General Hospital, <sup>10</sup>Stanford University Medical Center, <sup>11</sup>Washington University, <sup>12</sup>Lenox Hill Heart and Vascular Institute of New York, <sup>13</sup> Cedar Sinai Medical Center, <sup>14</sup> Saint Vincent Hospital at Worcester Medical Center, <sup>15</sup>Miami Children's Hospital

Corresponding Author:

Carl L. Tommaso MD (Chair, Writing Committee)  
Director, Cardiac Catheterization Laboratory  
Skokie Hospital, NSUHS  
9669 N. Kenton Avenue  
Skokie, IL  
(p) 847 675-8990  
(f) 847 675-1664  
(e) ctommaso@northshore.org

This article has been copublished in Catheterization and Cardiovascular Interventions, The Journal of Thoracic and Cardiovascular Surgery, The Annals of Thoracic Surgery, and The Journal of the American College of Cardiology.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/ccd.24394

© 2012 by the Society for Cardiovascular Angiography and Interventions, The Journal of Thoracic and Cardiovascular Surgery, The Annals of Thoracic Surgery, and The Journal of the American College of Cardiology.

## **Preamble**

The granting of staff privileges to physicians is an important mechanism to ensure quality care. The Joint Commission on Accreditation of Healthcare Organizations requires that medical staff privileges be based on professional criteria specified in medical staff bylaws. Physicians are charged with defining the criteria that constitute professional competence and with evaluating their peers accordingly. With the evolution of TAVR, an important opportunity arises for both cardiologists and surgeons to come together to identify the criteria for performing these procedures. The Society for Cardiovascular Angiography and Interventions (SCAI), American Association for Thoracic Surgery (AATS), American College of Cardiology Foundation (ACCF), and the Society of Thoracic Surgeons (STS) have therefore joined together to provide recommendations for institutions to assess their potential for instituting and/or maintaining a transcatheter valve program. This paper concerns Transcatheter Aortic Valve Replacement (TAVR). Since TAVR is in its infancy, there are few data upon which to base this consensus statement. Therefore, many of these recommendations are based on expert consensus. As the procedures evolve, technology changes, experience grows, and more data is accumulated, there will certainly be a need to update this consensus statement. However, with the FDA having just approved the first generation of TAVR devices, the writing committee and participating societies believe that the recommendations listed in this report serve as an appropriate starting point. In some ways, these recommendations apply to institutions more than to individuals. Since there is a strong consensus that these new valve therapies are best performed using a team approach, these credentialing criteria may be best applied at the institutional level.

Partnering societies used the ACCF's policy on relationships with industry to author this document [1]. To avoid actual, potential, or perceived conflicts of interest that may arise as a

result of industry relationships or personal interests among the writing committee, all members of the writing committee, as well as peer reviewers of the document, were asked to disclose all current healthcare-related relationships, including those existing 12 months before initiation of the writing effort. A committee of interventional cardiologists and surgeons was formed to include a majority of members with no relevant relationships with industry (RWI) and other entities, and led by an interventional cardiology co-chair and a surgical co-chair with no relevant RWI. Authors with relevant RWI were not permitted to draft or vote on text or recommendations pertaining to their RWI. RWI were reviewed on all conference calls and updated as changes occurred. Author and peer reviewer RWI pertinent to this document are disclosed in Appendixes 1 and 2, respectively. In addition, to ensure complete transparency, authors' comprehensive disclosure information (including RWI not pertinent to this document) is available as an online supplement to this document. The work of the writing committee was supported exclusively by the partnering societies without commercial support. Writing committee members volunteered their time to this effort. Conference calls of the writing committee were confidential and attended only by committee members.

SCAI, AATS, ACCF, and STS believe that adherence to these recommendations will maximize the chances that these therapies will become a successful part of the armamentarium for treating valvular heart disease in the United States. In addition, these recommendations will hopefully facilitate optimum quality during the delivery of this therapy, which will be important to the development and successful implementation of future, less invasive approaches to structural heart disease.

## **Introduction**

Enabled by the development of new technologies, treatment of valvular heart disease by transcatheter techniques is becoming a favored approach of cardiac providers, resulting in less invasive treatment for patients previously treatable only with open-heart surgery or, in many cases, not treatable at all. Recognition from the medical community of the applicability, effectiveness and practicality of catheter-based transcatheter valve therapies has further increased interest in these treatments. Training program content, standards, credentialing, and board certifications for cardiac surgical procedures and percutaneous coronary intervention are well developed, but no such structure exists in the field of percutaneous structural or valvular heart disease therapies. The purpose of this paper is to outline criteria for operator and institutional requirements in order to enable institutions and providers to participate responsibly in this new and rapidly developing field.

The emergence of transcatheter aortic valve repair and implantation as an alternative to traditional surgical therapy for valvular diseases has been facilitated by innovative devices, rapidly developing techniques, and careful patient selection. The combination of interventional skills, equipment, collaborative clinical management, surgical approaches, techniques, and decision-making distinguish the qualifications to participate in this field as unique, as does the complexity of the patients requiring these therapies [2],[3]. Given both the high-risk nature of these catheter interventions and the availability of established alternative treatment options using traditional surgical approaches, several considerations are important for institutions and operators planning to implement these new technologies.

Defining operator and institutional requirements for these novel therapies is an important first step in ensuring their optimal implementation.

Establishing a structural heart disease intervention therapy program requires several key components (Table 1). The defining principle is that this effort is a joint, institutionally-based activity for cardiologists and cardiac surgeons [4]. Thus, the specialty that provides some of these components will vary from program to program. A TAVR program that uses only one specialty is fundamentally deficient, and valve therapy programs should not be established without this multidisciplinary partnership. Comprehensive multidisciplinary teams (MDT) are, therefore, required for transcatheter valve therapies.

#### KNOWLEDGE BASE AND SKILLS

The critical cornerstone for establishing a transcatheter valve program is the formal collaborative effort between interventional cardiologists and cardiac surgeons. This element is essential for establishing a transcatheter valve program. No one individual, group, or specialty possesses all the necessary skills for best patient outcomes [5]. The over-arching goal of these programs must be to provide the best possible patient-centered care [6].

Since these are new techniques, the correlation between operator experience and performance metrics for these procedures has yet to be established. The current pool of trained individuals is comprised predominantly of those who have participated in industry-sponsored trials aimed at device approval. Therefore, the translation of currently available experiences with transcatheter valve therapies to the “real world” has yet to be evaluated in the United States.

Several core concepts should be implemented for all physicians performing these procedures, irrespective of their specialty background [7],[8]. They should all possess extensive knowledge of valvular heart disease, including the natural history of the disease, hemodynamics, appropriate diagnostics, optimal medical therapy, application and outcome of invasive therapies, and procedural and perioperative care [9].

The ability to interpret echocardiographic and other radiographic images (obtained at baseline, during the procedure, and follow-up) is critically important. MDTs and procedural teams need to possess echocardiographic interpretation skills for transthoracic and transesophageal studies. The use of 3D and 4D echoes may evolve to become essential diagnostic tools. Expertise in the interpretation of CT scans of the iliofemoral vessels, cardiac anatomy, as well as aortic valvular anatomy is critical for determining patient eligibility, and the approach for procedures [8],[10].

As noted, there is as yet no demonstrated direct correlation between operator experience with specific procedures and the skills necessary to perform transcatheter valve procedures, although there are some procedures that require similar prerequisite skills such as balloon aortic valvuloplasty for TAVR. There are, however, some core concepts that professional societies have accepted as important for both facilities and operators [11],[12]. Minimum training for specific procedures and devices will, for the immediate future, be primarily dictated by Food & Drug Administration (FDA) approval requirements. Simulation is likely to play a significant role in technical training and proficiency maintenance for these evolving procedures [13],[14],[15],[16],[17]. As these procedures become integrated into mainstream care delivery, the strategy for training will likely need to be revised.

Minimum requirements for transcatheter valve therapies include an understanding of basic radiation safety necessary for optimal imaging, operator and patient exposure protection, and knowledge of the use of x-ray contrast agents, which may not be standard in cardiac surgery training and experience.

Training in the use of closed systems for hemodynamic monitoring and contrast injections will result in optimal integration into catheterization laboratories and hybrid environments. Catheter and wire skills, including knowledge of the use of various techniques and the equipment available to access complex anatomy and negotiating necessary vascular and anatomic structures are required. Understanding of the interplay of wires, catheters, and anatomy is required for completion of these procedures. These skills can be acquired in a variety of ways. Prior experience with a variety of interventional techniques is important. These include but are not limited to:

- Coronary diagnostic procedures
- Coronary interventions
- Peripheral vascular diagnostic procedures
- Peripheral vascular interventions
- Balloon aortic, mitral and pulmonic valve dilatation
- Stent implantation in right ventricle outflow tract and pulmonary arteries
- intraaortic balloon pump (IABP), other cardiac support device placement, including initiation of percutaneous cardiopulmonary bypass
- percutaneous ventricular assist device (pVAD) placement
- endovascular aneurysm repair (EVAR) or thoracic endovascular aortic repair (TEVAR) procedures
- Transseptal techniques
- Coronary sinus access
- Large vessel access and closure

Operators should also have experience with specific catheter-based techniques required for valve interventions. Similarly, surgeons should have experience with transapical approaches for left

ventricular assist device (LVAD) placement and care of similar high-risk patients in order to perform transapical transcatheter aortic valve replacement (TAVR) [11, 12],[18]. The experience of an interventionalist or surgeon should be relevant to the transcatheter valve procedure undertaken. In this document, attention will focus upon cardiac surgery and interventional cardiology experience relevant to aortic valve intervention

The concept of sterile technique must be completely understood and stringently applied to the delivery of transcatheter valve therapies. Interventionalists must understand and be able to function in an environment that has more stringent sterile technique requirements than are common in catheterization laboratories. As one of the leaders of the team performing these procedures, the interventionalist must be able to enforce compliance with these standards. These procedures may involve open or partially open surgical components. Also, large devices that possess the same risk of infection as conventional valve prostheses will be implanted, especially for valve replacement procedures. Operating theater standards for sterile technique are therefore mandatory in order to ensure best patient outcomes.

## FACILITIES

The institution should have an active valvular heart disease surgical program with at least two institutionally-based cardiac surgeons experienced in valvular surgery, and should contain a full range of diagnostic imaging and therapeutic facilities including:

1. Cardiac catheterization laboratory or hybrid operating room/cath lab equipped with a fixed radiographic imaging system with flat-panel fluoroscopy, offering catheterization laboratory quality imaging. A bi-plane unit may be advantageous, particularly for congenital heart disease.
2. Non-invasive imaging

- a. Echocardiographic laboratory. Transthoracic and transesophageal echocardiographic capabilities with sonographers and echocardiographers experienced in valvular heart disease. Access to 3D echocardiography is preferable.
  - b. Vascular laboratory (non-invasive) with vascular specialists capable of performing and interpreting vascular studies
  - c. Computed tomography (CT) laboratory with CT technologists and specialists who can acquire and interpret cardiac CT studies
3. Physical space – The implantation suite must have a sterile environment that meets operating room standards. Furthermore, it must have sufficient space to accommodate the necessary equipment for uncomplicated implantations as well as any additional equipment that may be necessary in the event of complications. This includes space for anesthesiology, echocardiography, and cardiopulmonary bypass equipment and personnel. A specifically designed hybrid operating room (OR) interventional suite is ideal; however, in the absence of such a facility, the interventional cardiac suite should have:
- a. Circulating heating, ventilation, and air conditioning (HVAC) laminar flow diffusers (providing smooth, undisturbed air flow and usually placed directly over the procedure table) to meet air requirements for surgery rooms
  - b. Asymmetrical/symmetrical 6-lamp 2 x 4 troffers (the inverted, usually metal trough suspended from the ceiling as a fixture for fluorescent lighting) to provide adequate high-output lighting for surgical intervention
  - c. Adequate number of power receptacles that meet surgical equipment requirements

- d. Capability of running cardio-pulmonary bypass apparatus in the interventional suite
- e. Gas outlets for the anesthesia machine
- f. Adequate room size to accommodate the standard equipment required in a cardiac catheterization laboratory (e.g., High-Definition (HD) displays and monitors, O<sub>2</sub> analyzer, defibrillator/resuscitation cart, O<sub>2</sub> supply, suction, compressed air, CO-oximeter, activated clotting time (ACT) analyzer)
- g. Minimum room size of 800 square feet (74.3m<sup>2</sup>) to accommodate echocardiographic equipment, sonographers, anesthesia equipment, emergency CT surgical team and cardiopulmonary bypass equipment (e.g., surgeon, assistant, scrub tech, pump techs), if needed
4. Fungible equipment – The interventional suite should stock a large variety of fungible equipment, including various access kits, endovascular sheath and introducers ranging from 4 F to 26 F in various lengths, a wide range of guide wires for various purposes, cardiac diagnostic and interventional catheters, vascular closure devices, balloon dilatation catheters ranging from 2 mm to 30 mm in diameter and of various lengths and profiles, bare metal and covered stents (e.g., coronary and peripheral), occlusive vascular devices, snares and other retrieval devices, drainage catheters, and various implantable device sizes with their delivery systems
5. Post-procedure intensive care facility with personnel experienced in managing patients who have undergone conventional open-heart valve procedures
6. Use of mobile C-arm imaging system in an operating room is not adequate

7. HYBRID OR - The soon to be published *ACCF/SCAI 2012 Clinical Expert Consensus*

*Document on Cardiac Catheterization Laboratory Standards Update* will outline the specifications for a hybrid Cath Lab/OR. Though this is preferable, it is not a prerequisite since it is not available at many institutions

Most importantly, there must be dedication on the part of the hospital to provide these services and support, both financially and with no time constraints on the personnel involved. A dedicated administrator as a member of the team is necessary.

OTHER INSTITUTIONAL RESOURCES - Facilities, equipment (disposables and capital equipment), personnel, support, and commitment

For pre- and post-procedure care and joint formal multidisciplinary patient consultation, adequate outpatient clinical care facilities are necessary. Appropriate office space for the medical, nursing, and technical personnel involved is also required, preferably in a central setting. Ancillary testing facilities (i.e., pulmonary function, echocardiography, vascular Duplex scanning, clinical laboratory, multi-slice computed tomography (MSCT)) should be of high quality and able to accommodate the patient load on a timely basis.

By their very nature, these complex procedures should only be performed in institutions that currently and routinely perform large volumes of surgical aortic valve operations with outcomes that equal or exceed those established nationally for similar procedures. Similarly, only institutions with interventional cardiology programs that have established and successful programs with balloon aortic valvuloplasty, catheter closure of periprosthetic valvular leaks, insertion of ventricular septal closure devices with outcomes that equal or exceed those established nationally for similar procedures should develop an integrated structural heart MDT.

The institutional commitment required for a successful program goes beyond the necessary space, personnel, and specialized facilities set forth above. The complex and time consuming pre-procedure patient triage process and the amount and intensity of post-procedure patient care after discharge are very labor intensive for the physician and nursing staff, as are the informed consents and communications with patients, families and referring providers. In addition to supporting the core nursing and technical support staff, arrangements between the institution and the physicians need to be structured to reimburse physician efforts dedicated to non-reimbursable hours of clinical care and medical management of the program.

The complexity of transcatheter valve procedures and the magnitude of institutional resources required are similar to established heart transplant and cardiac assist device programs, where dedicated professionals, a minimum of infrastructure, MDT, registered nurse/nurse practitioner (RN/NP), providers, coordinators, databases and quality reporting are essential for optimal patient outcomes. This concept was endorsed by the Centers for Medicare and Medicaid Services (CMS) through the establishment of certification criteria for the use of heart transplantation and cardiac assist devices in centers and, moreover, for eligibility for reimbursement of services provided. The same regulatory system was applied to professionals providing these services. Transcatheter valve treatment programs should undergo a similar regulatory process with CMS endorsement. Centers should be approved for transcatheter aortic valve programs based on a minimum number of cases per year, and perioperative and 1-year outcomes above a minimum threshold.

Long-term outcome reporting is obligatory, in order to track not only survival, but also other parameters including periprocedural complications (CVA, vascular, renal, infectious, etc.), aortic regurgitation, the need for re-intervention, subsequent surgery and quality of life. This type of

reporting is essential, since long-term outcome goals for these new procedures have not been established at this early stage. Participation in a national data registry (e.g. STS/ACC TVT Registry) is mandatory.

#### MULTIDISCIPLINARY TEAM

The use of a team approach has been shown to improve outcomes in these types of complex procedures[19]. The MDT necessary for a TAVR program is highlighted by the collaboration between the interventional cardiologist and cardiac surgeon. The MDT, however, goes well beyond this collaboration, and must include key providers from other physician groups (e.g. Anesthesiology, Radiology, Non-invasive Cardiology, Intensive Care, etc.). In addition to the individual physicians, other components that extend to various departments are necessary. The idea that the MDT is comprised of individual physicians working in a room performing the procedures is a superficial view that does not take into account the level of resources necessary for a successful valve therapy program. The interaction among specialists in the MDT is fundamental, particularly for pre-procedure patient evaluation and selection. It is equally fundamental that the patient be at the very center of all discussions and decision-making regarding the best therapy in her/his particular circumstance. While there is great excitement about the application of transcatheter valve therapies, most of these therapies will only be indicated for a small portion of the population for the immediate future. Proper decision-making and determination of best options for any given patient require an evaluation by the MDT [20].

On-site valve surgery is an essential component of any valve therapy program. The requirement for on-site valve surgery is based not only on the potential need for emergency or “back-up” surgery for percutaneous patients, but more importantly on the quality of patient evaluation and selection, decision-making, intra-procedure management, and post-procedure care and outcomes.

A cardiac surgeon and an interventional cardiologist must evaluate every case. Interplay between interventional cardiologists and cardiac surgeons represent only part of the benefit of the MDT.

As noted above, additional critical contributions are provided by cardiac anesthesiology, imaging specialists in both cardiology and radiology, and by the many people who extend beyond the physician members of the team. The MDT is led by a core of physicians from interventional cardiology, cardiac surgery, cardiac anesthesiology and intensive care and cardiac imaging departments, along with congenital heart disease specialists and surgeons, in some instances. Depending on the institutional organization and the needs of the patient, vascular surgery and interventional radiology departments may also participate in the MDT. Additional team members include nurse practitioners from all of these fields, research coordinators, and a dedicated administrator.

The function of the MDT is essential in pre-procedure patient selection, intra-procedure management and problem solving, post-procedure management, post-discharge follow-up, and outcome studies. During procedures, emergencies or unanticipated needs may arise as a matter of course even in the most straightforward situation. The immediate availability of MDT support to help with decision-making or with therapy is essential. A clear definition of roles for different specialties as well as effective communication, which may be different from that of conventional procedures, is critical for successful outcomes. Difficult post-procedure courses are common in the high-risk patients who comprise a large part of the target population for both transcatheter and operative valve therapies. A team approach to problem solving in this setting is critical.

Another important part of patient management is the familiarity that the intensive care unit and the monitored step-down floor staff have with the specific details of each form of valve therapy.

After the post-procedure management phase, long-term, post-hospital follow-up for this select

group of patients is also part of the MDT's responsibility. Planning for and resourcing this important phase of care is incumbent upon the MDT. Post approval registries (e.g., STS/ACC TVT Registry) will be required for many of the new transcatheter valve therapies, and, therefore, a data collection/research unit within the MDT is another required component.

For sites with no prior trial experience in aortic transcatheter therapies, background experience from related procedures is important. The surgical use of ventricular assist device (VAD) support or apical conduit therapy for aortic stenosis or left ventricular apical venting during aortic aneurysm procedures provides excellent experience for management of apical access for TAVR. For transcatheter procedures that do not directly involve the surgeon as a procedure operator, the role of the cardiac surgeon remains critically important. The surgeon has many roles and is often a patient advocate and/or referring physician, and is a necessary scientific study participant in all of these device applications. The surgeon is familiar with established standards of surgical care for application in transcatheter therapies and is frequently in charge of assessing high-risk patients for catheter-based therapy as an alternative to surgery. In a valve therapy program, neither the surgeon nor the cardiologist should be in charge of the assessment, but rather the MDT.

**IN ALL TRANSCATHETER AORTIC PROCEDURES, THE INTERVENTIONALIST AND THE SURGEON SHOULD BE PRESENT FOR THE CRITICAL PORTIONS OF THE PROCEDURE.**

Another mechanism for promoting a team approach that involves both surgeons and cardiologists is split or shared physician reimbursement for these procedures, which this writing group strongly endorses. This important principle will ensure that surgeons and cardiologists participate jointly in performing procedures, and that each patient receives the best and most patient-centered treatment.

The MDT should meet formally as a group on a regular basis (aside from the usual “cath conference”) to review all patients referred for procedures, performance of recent procedures (to discuss both favorable and unfavorable outcomes), and follow-up of prior procedures.

#### FUNCTION OF THE MULTI-DISCIPLINARY TEAM

Programmatic success depends on the ability of the MDT to function effectively in the best interest of a given patient. To do so, the MDT must work cohesively through the processes of patient selection, procedural planning, procedural conduct, periprocedural care and longitudinal follow-up. Through each phase of this continuum, the individual skills of the MDT members should be brought to bear upon the process.

The procedural success of transcatheter valve therapies begins with patient selection. Given the complexity of the decision-making process surrounding these procedures, it is necessary that all MDT members provide objective input and judgments from the outset of a patient evaluation. In order that true informed consent be obtained, the patient must remain at the center of the deliberations of the MDT at all times, and must be involved in the discussions regarding her/his therapeutic approach and goals. The patient selection process may be initiated by use of regularly scheduled patient selection conferences attended by all MDT members. Such conferences are analogous to transplant patient selection committee meetings, and provide a venue in which patient-specific data and imaging are formally presented and discussed by the MDT. The respective expertise of each discipline represented among MDT members may then be synthesized into a patient-specific recommendation.

Direct patient evaluation by cardiologists and cardiac surgeons may be accomplished jointly and, if possible, simultaneously in a venue such as a multi-disciplinary valve program clinic. Not only does such a clinic provide convenience for the many patients who are elderly and fragile, but it also provides an opportunity for cardiac surgeons and cardiologists to jointly examine and evaluate complex patients.

In so doing, the expertise and judgment of both disciplines may be woven into a patient-specific decision. The participation of anesthesiologists in these clinics may also be useful.

Following the decision that a given patient is an appropriate candidate for TAVR, the procedure must then be carefully planned. Cardiac surgical teams are familiar with, and routinely use the concept of “pre-procedure briefings” prior to complex cardiac surgical operations. In such briefings, all team members (i.e., surgeons, anesthesiologists, perfusionists, nurses, technicians) discuss the intended procedure, including the steps of the planned procedure, the specific tools and equipment needed (beyond those typically used), the possible complications that may arise during the course of the procedure, and the contingency plans that will be implemented should the unexpected occur. All members of the team may then initiate the planned procedure with a common understanding of its conduct and what will happen if the plan needs to change.

As integral members of the MDT, the cardiologist, cardiovascular surgeon and the catheterization team will participate actively in this pre-procedure planning and MDT briefing, which is so important for the procedural success of transcatheter valve therapies. During the procedure, emergency situations and unexpected needs may arise. The immediate availability of MDT physician support in emergency decision-making and therapy is essential. It is therefore

important that the roles of the various specialties be clearly delineated during pre-procedure planning.

Patients that undergo transcatheter valve therapies are often elderly and frail with multiple comorbidities. Post-procedure care of such patients may be difficult and entail the management of multiple organ system dysfunctions. In many cases, the initial post-procedure care should be provided in an intensive care setting. A team approach to the care of these patients and to problem solving, is important and should include physicians skilled in critical care medicine.

Once in-patients are able to leave the intensive care environment, they should be attended by a unit specializing in the care of patients with cardiac diseases, and this unit should be equipped with telemetry-monitored beds. Again, a team approach is important for success. The team of physicians, nurses, occupational and physical therapists, and other members must have an understanding of the pathophysiology of the particular valve condition, as well as the nuances of care for patients who have undergone cardiac surgery and interventional cardiology procedures.

Procedural success of transcatheter valve therapies must be determined via longitudinal outcomes. Long-term follow-up of these patients is an important element of the MDT approach. Post-FDA approval registries (e.g., STS/ACC TVT Registry) will be required for most transcatheter valve therapies. Therefore, a long-term relationship between the patient and the MDT must be established in order to undertake the needed alterations in medical therapy, serial echocardiographic imaging, and monitoring of devices. Likewise, changes in patient functional status, heart failure class, potential device-related complications, and other such conditions must be carefully tracked. A valve program clinic can provide a venue for this type of long-term follow-up.

The post-market surveillance of transcatheter valve devices will be an extremely important function of the MDT. Participation in device-specific registries (e.g., STS/ACC TVT Registry) can be challenging and requires an institutional infrastructure and commitment that includes experienced data managers with a background in cardiac disease, funding, office space, and computer resources. It requires a clinical research unit with rigorous attention to detail, and the collection of accurate data as an integral part of the function of the MDT.

CRITERIA FOR ESTABLISHING A TRANSCATHETER VALVE PROGRAM AND MAINTENANCE OF COMPETENCE

An important issue in the establishment of a transcatheter valve program is the clinical or referral base for ensuring an adequate number of patients to provide for the viability of a program. The following requirements are important for the establishment of a successful trans-catheter valve program:

TABLE 1	<b>TRANSCATHETER AORTIC VALVE REPLACEMENT</b>
<b>INSTITUTIONAL INTERVENTIONAL PROGRAM</b>	1000 CATH/400 PCI PER YEAR*
<b>TAVR INTERVENTIONALIST</b>	<p>100 STRUCTURAL PROCEDURES LIFETIME OR 30 LEFT SIDED STRUCTURAL PER YEAR OF WHICH 60% SHOULD BE BALLOON AORTIC VALVULOPLASTY</p> <p>(LEFT SIDED PROCEDURES INCLUDE EVAR, TEVAR, BALLOON AORTIC VALVE (BAV), AORTIC VALVE (AV) AND MITRAL VALVE (MV) PROSTHETIC LEAK CLOSURES AND VSD CLOSURES. (ATRIAL SEPTAL DEFECT/PATENT FORMAN OVALE (ASD/PFO) CLOSURE ARE NOT CONSIDERED LEFT SIDED PROCEDURES</p>
	SUITABLE TRAINING ON DEVICES TO BE USED

<b>INSTITUTIONAL SURGICAL PROGRAM</b>	<p>50 TOTAL AVR PER YEAR OF WHICH AT LEAST 10 AORTIC VALVE REPLACEMENT (AVR) SHOULD BE HIGH-RISK (STS SCORE <math>\geq</math> 6)</p> <p>MINIMUM OF 2 INSTITUTIONALLY-BASED CARDIAC SURGEONS IN PROGRAM (MORE THAN 50% TIME AT HOSPITAL WITH SURGICAL PROGRAM)</p>
<b>TAVR SURGEON</b>	<p>100 AVR CAREER, AT LEAST 10 OF WHICH ARE “HIGH-RISK” (STS SCORE <math>\geq</math> 6) OR 25 AVR PER YEAR OR 50 AVR IN 2 YEARS AND AT LEAST 20 AVR IN LAST YEAR PRIOR TO TAVR INITIATION</p>
	<p>EXPERIENCE WITH, AND MANAGEMENT OF PERIPHERAL CARDIOPULMONARY BYPASS EXPERIENCE WITH OPEN RETROPERITONEAL EXPOSURE OF, AND SURGICAL INTERVENTION ON THE ILIAC ARTERIES</p>
	<p>ALL CASES MUST BE SUBMITTED TO A SINGLE NATIONAL DATABASE</p>
<b>EXISTING PROGRAMS</b>	<p>30 TAVR (TOTAL EXPERIENCE)</p>
	<p>ONGOING CONTINUING MEDICAL EDUCATION (CME) (OR NURSING/TECHNOLOGIST EQUIVALENT) OF 10 HOURS PER YEAR OF RELEVANT MATERIAL</p>
	<p>ALL CASES MUST BE SUBMITTED TO A SINGLE NATIONAL DATABASE</p>

<p>TRAINING</p>	<p>CARDIOLOGISTS MUST BE BOARD CERTIFIED/ELIGIBLE IN INTERVENTIONAL CARDIOLOGY</p> <p>SURGEONS MUST BE BOARD CERTIFIED /ELIGIBLE IN THORACIC SURGERY</p> <p>ADDITIONAL OPERATORS WHO ARE TRAINED OR EXPERIENCED IN STRUCTURAL HEART DISEASE AND HAVE UNRESTRICTED HOSPITAL PRIVILEGES IN STRUCTURAL PROCEDURES, MAY ALSO BE PART OF THE INTERVENTIONAL OPERATING TEAM WITH THE INTERVENTIONAL CARDIOLOGIST AND CARDIOVASCULAR SURGEON</p>
<p><b>NEW PROGRAMS</b></p>	<p>20 TAVR PROCEDUES/ YEAR or 40 TAVR PROCEDURES OVER 2 YEARS</p> <p>30-DAY ALL-CAUSE MORTALITY &lt;15%</p> <p>30-DAY ALL-CAUSE NEUROLOGIC EVENTS INCLUDING TRANSIENT ISCHEMIC ATTACK (TIA'S)&lt;15%</p> <p>MAJOR VASCULAR COMPLICATION &lt;15%**</p> <p>&gt;90% INSTITUTIONAL FOLLOW-UP</p> <p>60% 1-YEAR SURVIVAL RATE FOR NON-OPERABLE PATIENTS (COHORT B) – AFTER THE PROGRAM HAS BEEN RUNNING FOR 2 YEARS ( 2-YEAR AVERAGE)</p>
	<p>ONGOING CME (OR NURSING/TECHNOLOGIST EQUIVALENT) OF 10 HOURS PER YEAR OF RELEVANT MATERIAL</p>
	<p>ALL CASES MUST BE SUBMITTED TO A SINGLE NATIONAL DATABASE</p>

TRAINING	<ol style="list-style-type: none"><li>1) CARDIOLOGISTS MUST BE BOARD CERTIFIED/ELIGIBLE IN INTERVENTIONAL CARDIOLOGY</li><li>2) SURGEONS MUST BE BOARD CERTIFIED /ELIGIBLE IN THORACIC SURGERY</li><li>3) ADDITIONAL OPERATORS WHO ARE TRAINED OR EXPERIENCED IN STRUCTURAL HEART DISEASE AND HAVE UNRESTRICTED HOSPITAL PRIVILEGES IN STRUCTURAL PROCEDURES, MAY ALSO BE PART OF THE INTERVENTIONAL OPERATING TEAM WITH THE INTERVENTIONAL CARDIOLOGIST AND CARDIOVASCULAR SURGEON</li></ol>
----------	---

\* With acceptable outcomes for conventional procedures compared to NCDR benchmarks

\*\* According to BARC-2 definitions

Once chosen for participation as TAVR programs, either as existing programs, or as new programs, in order to maintain ongoing approval for participation, sites will be monitored to ensure that they continue to satisfy both the volume and outcomes criteria as described in Table 1.

Unlike the experience with percutaneous coronary intervention (PCI), where data attest to the relationship between the volume of procedures and outcomes, there are little or no data on which to draw conclusions as to the volume-outcome relationship for TAVR. Therefore, the above recommendations are constructed to: 1) ensure patient safety, 2) demonstrate that there is a commitment on the part of the institution to the structural heart disease program, and 3) use existing volume as a surrogate for an established valve program to ensure adequate patient volumes for the establishment of a sustainable and high-quality transcatheter valve program. As experience grows and more data become available, these recommendations will undoubtedly be refined.

## AORTIC VALVE REPLACEMENT

Surgical aortic valve replacement is the treatment of choice for patients with severe aortic valve disease [21]. However, a significant percentage of those patients are not offered the procedure or simply refuse to undergo it on the basis of excessive risks, both real and perceived[22]. It is in this context that the possibility of transcatheter aortic valve implantation has become a reality for patients outside of the United States. Furthermore, the cohort B results of the PARTNER (Placement of AoRticTraNscathetER) trial (non-operable patients) would indicate that medical therapy, including balloon aortic valvuloplasty (BAV), should be reserved only for patients who do not qualify for the procedure on the basis of their anatomy or clinical characteristics[23].

Considering the reports of successful transcatheter treatment of aortic valve stenosis using balloons [24] and subsequent reports of poor long-term outcomes due to early restenosis [25], the idea of developing a transcatheter aortic valve was a logical progression. Initial animal studies [26] and subsequent human implantations have led to the progressive development of this technology.

Results from the PARTNER trial using the Edwards Sapien Valve (Edwards Lifesciences, Irvine, California), the first randomized trial of this technology, have established its place as a treatment of severe symptomatic aortic stenosis [23],[27]. The trial consisted of two arms: cohort A (n=699), the high risk surgical group, which randomized patients to surgical aortic valve replacement or TAVR, and cohort B (n=358), the non-surgical group randomized to medical therapy, which could include valvuloplasty or TAVR. Results from the non-inferiority cohort A group showed a one year all-cause mortality of 26.8% in the surgical arm and 24.2% in the TAVR arm, a hazard ratio (HR)(95% CI) of 0.93 (0.7, 1.22), which met the non-inferiority endpoint,  $p=0.001$ (for a non-inferiority margin of 7.5 percentage points). There were differences in outcomes between the groups, the most worrisome being a higher incidence of stroke or TIA at 30 days and at one year in the TAVR arm (at 30 days: 5.5% vs. 2.4%,  $p=0.04$  and at 1 year: 8.3% vs. 4.3%,  $p=0.04$ ). The need for a new permanent pacemaker was not different between the groups, 5.7% in the surgical arm vs. 5.0% in the TAVR arm,  $p=0.68$ . In cohort B, TAVR was found to be superior to medical therapy with an all-cause mortality at one year of 50.7% in the medical arm vs. 30.7% in the TAVR arm, HR (95% CI) 0.54 (0.58, 0.78,  $p<0.001$ ). These findings suggest that it is necessary to treat 5 patients to prevent 1 death[28]. PARTNER II, which uses a new, lower profile delivery system, is in the early stages of patient recruitment. To

date, there are no randomized data regarding the CoreValve prosthesis (Medtronic, Minneapolis, MN) but it has received the CE mark in Europe, as has the Sapien valve.

Registry data appear to show favorable hemodynamic outcomes and acceptable mortality [29], [30]. In this large multicenter registry consisting of 14 centers reporting on 663 consecutive patients, [30] procedure success was 98% with a procedural mortality of 0.9%. Thirty-day mortality was 5.4% and 15.0% at one year. Stroke during the procedure and at one year was 1.2% and 2.5%, respectively. The requirement for a permanent pacemaker was 19.1% at one year, a rate similar to other registries reporting outcomes using this valve. Independent predictors of mortality at 30 days included conversion to open heart surgery (odds ratio [OR] 38.68), cardiac tamponade (OR 10.97), major access site complications (OR 8.47), left ventricular ejection fraction <40% (OR 3.51), prior balloon valvuloplasty (OR 2.87), and diabetes mellitus (OR 2.66). Predictors of late mortality (30 days to one year) included prior stroke (HR 5.47), post procedural paravalvular leak  $\geq 2+$  (HR 3.79), prior acute pulmonary edema (HR 2.70), and chronic kidney disease (HR 2.53). The CoreValve US Pivotal Trial is currently randomizing patients in a similar fashion to the PARTNER trial and should yield additional important data on this valve.

#### RESEARCH / REGISTRY

FDA clearance of a novel valve repair or replacement prosthesis does not guarantee that the device will continue to demonstrate long-term efficacy equal to currently available options, or that its application will be limited to the initially approved patient subsets. Post-market studies organized through individual institutions or multicenter study groups and registries managed by industry and professional societies are essential for ensuring continued short-term safety, and for

determining long-term efficacy. Only with such data can we consider application of new valve prostheses to a wider patient population outside the boundaries of the study groups examined during FDA trials. Centers that incorporate transcatheter-based therapies into their practice absolutely must participate in a cardiac surgery or cardiology national database such as the Society of Thoracic Surgeons' National Database, Northern New England Cardiac Disease Study Group, American College of Cardiology's National Cardiac Data Registry, or an equivalent database [31],[32],[33]. These databases facilitate continued analysis of early outcomes on a national level; however, most do not permit analysis of results beyond the 30-day window, mandating the development of implant registries with patient consent, permitting late follow-up for survival and valve-related complications and re-interventions. These valve repair and replacement procedures should be registered in databases capable of providing both acute outcome and long-term follow-up data.

Early post-procedure morbidity and mortality analyses, while important for initial and continued implant safety assessment, are not sufficient to evaluate the efficacy of valve repair or replacement prostheses. Studies on long-term follow-up survival and, more importantly, structural valve degeneration, and the need for reintervention, are essential. Recent attempts to link the STS National Database to an administrative survival database (Social Security Death Master File) or to CMS Medicare data have been promising however, survival data alone will not be sufficient for transcatheter valve registries (e.g., STS/ACC TVT Registry) [34]. Risk adjustment using only administrative data is challenging and may be more important in this highly complex patient population. Transcatheter valve registries must incorporate late assessment of structural valve degeneration and the need for late reoperation or reintervention. Long-term function data is also essential before application of new valve repair or replacement

technology can be considered for lower risk and younger patients. Clinicians must be careful not to extrapolate outcomes generated during Food and Drug Administration (FDA) trials to patients that do not reasonably approximate the trial study populations. At present, it is not acceptable for clinicians to apply “off label” transcatheter techniques to patients who are otherwise excellent candidates for conventional valve repair or replacement outside the confines of a randomized, controlled trial or, at a minimum, multicenter, national, prospective studies.

The potential negative impact of a valve prosthesis recall, in regards to the increased risk associated with reoperative surgery for a permanent implant, far outweighs that associated with the removal of a pharmacologic agent from the market [35]. Thus, the allocation of adequate funding to allow complete follow-up studies is essential, including financial support from industry, the FDA, National Institutes of Health (NIH), CMS, and professional societies, with scientific oversight distanced from industry and potential conflicts of interest. Post-market surveillance governed solely by industry self-regulation can be of dubious value. In a 2005 editorial, Eugene Blackstone cautioned, “Industry has not developed a viable mechanism for dealing with ‘bad news’, the disclosure of which often leads to the demise of the company” [36]. The Interagency Registry for Mechanically Assisted Circulatory Support (INTERMACS) exemplifies a potentially efficient model that could be emulated by professional societies for monitoring transcatheter inventions during the next decade, during which durability data will be generated [37].

Transcatheter valve repair or replacement devices are unique in that an understanding not only of early risk, but also of long-term durability, is essential for determining the appropriate patient subgroups for these therapies. In our opinion, it is the responsibility of professional societies to ensure adequate long-term data monitoring and provide oversight and guidance to industry on

the expectations for continued monitoring beyond the FDA approval phase of device development and implementation. Individual centers are also responsible for critically evaluating their own experience, through local and regional quality improvement initiatives, and for participating in national databases and registries (e.g., STS/ACC TVT Registry) that facilitate continued safety and efficacy in the assessment of novel, and, as yet, unproven, therapeutic options. Components of a national/international registry should include preoperative risk factors and valve assessment, intraoperative details, early postoperative morbidity, and late follow-up including survival, need for re-intervention, functional class, device-related complications and late assessment of valve performance. It is inappropriate to perform these novel and innovative procedures without the institutional infrastructure to ensure adequate early data collection and later follow-up.

**APPENDIX 1. Author Relationships With Industry and Other Entities (Relevant)—2012 Multisociety (AATS, ACCF, SCAI & STS) Writing Committee on Operator and Institutional Requirements for Transcatheter Valve Repair and Replacement**

<b>Committee Member</b>	<b>Employment</b>	<b>Consultant</b>	<b>Speaker's Bureau</b>	<b>Ownership/ Partnership / Principal</b>	<b>Personal Research</b>	<b>Institutional, Organizational or Other Financial Benefit</b>	<b>Expert Witness</b>
Michael A. Acker	Hospital of the University of Pennsylvania—Chief, Division of Cardiovascular Surgery	None	None	None	None	None	None
Gabriel S. Aldea	University of Washington Medical—Associate Professor	None	None	None	None	None	None
Joseph Bavaria*	Hospital of the University of Pennsylvania Health System Cardiothoracic Surgery—Vice-Chief, Division of Cardiothoracic Surgery; Brooke Roberts-William M. Measey Professor of Surgery; Director, Thoracic Aortic Surgery Program	St. Jude Medical	None	None	Medtronic Vascular	Edwards Lifesciences† Medtronic†	None
R. Morton (Chip) Bolman	Harvard Medical School—Chief, Division of Cardiac Surgery; Professor of Surgery	None	None	None	None	None	None

Duke Cameron	The Johns Hopkins Hospital	None	None	None	None	None	None
Larry Dean*	University of Washington School of Medicine—Professor of Medicine and Surgery	Emageon Philips Medical	None	None	Edwards Lifesciences‡	None	None
Ted Feldman*	Evanston Northwestern Healthcare Cardiology Division—Director, Cardiac Catheterization Laboratory	Abbott Boston Scientific Edwards Lifesciences WL Gore	None	None	Abbott Boston Scientific† Edwards Lifesciences WL Gore	None	None
David Fullerton	University of Colorado School of Medicine—Chief, Cardiothoracic Surgery	None	None	None	None	None	None
Ziyad M. Hijazi*	Rush University Medical Center—Professor, Pediatrics and Internal Medicine; Director for the Center of Congenital and Structural Heart Disease	Edwards LifeSciences‡ NuMED‡	None	Colibri Heart Valve†	None	None	None
Eric Horlick*	Toronto General Hospital—Asst. Professor of Medicine, University	Abbott Vascular AGA‡ Cordis Edwards	Biosense Webster	None	None	None	None

	Health Network	Lifesciences† Medtronic WL Gore					
D. Craig Miller*	Stanford University Medical School Dept. Cardiothoracic Surgery— Thelma and Henry Doelger Professor of Cardiovascular Surgery; Cardiovascular Surgical Physiology Research Laboratories— Director	Abbott Vascular MitraClip Medtronic Cardiovascular Division St. Jude Medical	None	None	Edwards Lifesciences (PARTNER Trial)†	Edwards Lifesciences (PARTNER Trial)†	None
Marc R. Moon	Barnes-Jewish Hospital—Joseph Bancroft Professor of Surgery, Division of Cardiothoracic Surgery; Program Director, Thoracic Surgery Residency Program	None	None	None	None	None	None

Richard R. Ringel	Johns Hopkins School of Medicine—Associate Professor of Pediatrics	None	None	None	None	None	None
Carlos E. Ruiz	Lenox Hill Heart and Vascular Institute of New York—Professor and Chief, Division of Pediatric Cardiology	None	None	None	None	None	None
Carl Tommaso	North Shore University HealthSystem—Medical Director of Cardiac Cath Lab	None	None	None	None	None	None
Alfred Trento	Cedar Sinai Medical Center—Director of Division of Cardiothoracic Surgery	None	None	None	None	None	None

Bonnie Weiner*	Saint Vincent Hospital at Worcester Medical Center/Fallon Clinic—Director of Interventional Cardiology Research; Imaging Core Lab Services—President ; Boston Biomedical Associates—Medical Director	LabCoat/Boston Scientific‡	Medtronic	None	None	Accreditation for Cardiovascular Excellence‡ Imaging Core Lab Services†	None
Evan M. Zahn	Miami Children's Hospital—Director of Cardiology	Medtronic† Edwards Lifesciences‡	None	None	None	None	None

This table represents the relationships of committee members with industry and other entities that were determined to be relevant to this document. These relationships were reviewed and updated in conjunction with all meetings and/or conference calls of the writing committee during the document development process. The table does not necessarily reflect relationships with industry at the time of publication. A person is deemed to have a significant interest in a business if the interest represents ownership of 5% or more of the voting stock or share of the business entity, or ownership of ≥\$10,000 of the fair market value of the business entity; or if funds received by the person from the business entity exceed 5% of the person's gross income for the previous year. Relationships that exist with no financial benefit are also included for the purpose of transparency. Relationships in this table are modest unless otherwise noted. Please refer to <http://www.cardiosource.org/Science-And-Quality/Practice-Guidelines-and-Quality-Standards/Relationships-With-Industry-Policy.aspx> for definitions of disclosure categories or additional information about the ACCF Disclosure Policy for Writing Committees.

According to the ACCF, a person has a **relevant** relationship IF: a) The relationship or interest relates to the same or similar subject matter, intellectual property or asset, topic, or issue addressed in the document; or b) The company/entity (with whom the relationship exists) makes a drug, drug class, or device addressed in the document, or makes a competing drug or device addressed in the document; or c) The person or a member of the person's household, has a reasonable potential for financial, professional or other personal gain or loss as a result of the issues/content addressed in the document.

\*Recused from writing initial text and voting on document recommendations due to relevant relationships with industry to this document.

†No financial benefit.

‡Significant relationship.

**APPENDIX 2. Reviewer Relationships with Industry and Other Entities (Relevant)—2012 Multisociety (AATS, ACCF, SCAI & STS) Writing Committee on Operator and Institutional Requirements for Transcatheter Valve Repair and Replacement**

<b>Peer Reviewer</b>	<b>Representation</b>	<b>Employment</b>	<b>Consultant</b>	<b>Speaker's Bureau</b>	<b>Ownership/ Partnership/ Principal</b>	<b>Personal Research</b>	<b>Institutional, Organizational, or Other Financial Benefit</b>	<b>Expert Witness</b>
Deepak L. Bhatt	Official Reviewer— ACCF Task Force on Clinical Expert Consensus Documents	VA Boston Health Care System—Chief, Division of Cardiology	None	None	None	<ul style="list-style-type: none"> <li>• Medtronic*</li> <li>• Ethicon*</li> </ul>	None	None
Ann Bolger	Content Reviewer— ACCF/AHA/ ACP Task Force on Clinical Competence and Training	University of California, San Francisco— Professor of Medicine	None	None	None	None	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Emmanouil Brilakis	Content Reviewer— ACCF Interventional Scientific Council	UT Southwestern Medical School— Director, Cardiac Catheterization Laboratory, VA North Texas Healthcare System	<ul style="list-style-type: none"> <li>St. Jude*</li> </ul>	None	None	None	<ul style="list-style-type: none"> <li>Abbott Vascular*</li> <li>Infraredx*</li> <li>Medtronic*</li> </ul>	None
John H. Calhoun	Official Reviewer— STS	University of Texas Health Sciences Center— Professor and Chair, Department of CT Surgery	None	None	None	None	None	None
George Dangas	Content Reviewer— ACCF Interventional Scientific Council	Mount Sinai Medical Center— Program Director, Interventional Cardiology	<ul style="list-style-type: none"> <li>Abbott</li> <li>Boston Scientific</li> <li>Johnson &amp; Johnson</li> <li>Medtronic</li> <li>St. Jude Medical</li> </ul>	None	None	None	<ul style="list-style-type: none"> <li>Abbott Vascular†</li> <li>Infraredx†</li> <li>Medtronic†</li> <li>W.L. Gore†</li> </ul>	Plaintiff, stroke, 2011

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Fred H. Edwards	Official Reviewer— STS	Shands Jacksonville – University of Florida— Professor of Surgery; Chief, Cardiothoracic Surgery	None	None	None	None	None	None
James I. Fann	Official Reviewer— AATS	Stanford University School of Medicine	None	None	None	• Medtronic†	None	None
James Fasules	Senior Vice President, Advocacy	American College of Cardiology	None	None	None	None	None	None
T. Bruce Ferguson	Content Reviewer— ACCF Interventional Scientific Council	East Carolina Heart Institute Brody School of Medicine at ECU— Professor of Surgery and Physiology	None	None	None	None	• Edwards Lifesciences	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Mario J. Garcia	Content Reviewer— ACCF Task Force on Clinical Expert Consensus Documents	Montefiore Medical Center-Albert Einstein College of Medicine— Chief, Division of Cardiology	<ul style="list-style-type: none"> <li>MD Imaging</li> </ul>	None	None	None	<ul style="list-style-type: none"> <li>Medtronic†</li> </ul>	None
Federico Gentile	Content Reviewer— ACCF Task Force on Clinical Expert Consensus Documents	Centro Medico Diagnostico	None	None	None	None	None	None
Leonard N. Girardi, MD	Official Reviewer— AATS	New York-Presbyterian/ Weill Cornell Medical College	None	None	None	None	None	None
Steve Goldberg	Official Reviewer— SCAI	University of Washington Medical Center	<ul style="list-style-type: none"> <li>AGA Medical</li> </ul>	None	None	None	<ul style="list-style-type: none"> <li>Amplatz Division of St. Jude Medical</li> </ul>	Plaintiff, Patent Litigation, 2010

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Frederick L. Grover	Content Reviewer—ACCF Surgeon Scientific Council	University of Colorado Department of Surgery—Professor and Chair	None	None	None	None	None	None
Robert Guyton	Content Reviewer—ACC Surgeon Scientific Council	Emory Clinic, Inc.—Professor and Chief, Division of Cardiothoracic Surgery	None	None	None	None	None	Defendant, Cardiac Surgery, 2011
Jonathan Halperin	Content Reviewer—ACCF/AHA/ACP Task Force on Clinical Competence and Training	Mount Sinai Medical Center—Professor of Medicine	<ul style="list-style-type: none"> <li>• Biotronix, Inc.*</li> <li>• Edwards Lifesciences</li> <li>• Johnson &amp; Johnson</li> <li>• Ortho-McNeil/Janssen Pharmaceuticals</li> </ul>	None	None	None	None	None
Howard Herrmann	Official Reviewer—SCAI	University of Pennsylvania Medical Center	<ul style="list-style-type: none"> <li>• W.L. Gore*</li> <li>• St. Jude Medical</li> </ul>	None	<ul style="list-style-type: none"> <li>• Micro Interventional†</li> </ul>	<ul style="list-style-type: none"> <li>• Edward Lifesciences*</li> <li>• W.L. Gore</li> </ul>	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Joerg Herrmann	Content Reviewer— ACCF Interventional Scientific Council	Mayo Clinic	None	None	None	None	None	None
David R. Holmes, Jr.	ACCF/AATS/ SCAI/STS TAVR ECD Writing Committee	Mayo Clinic— Consultant, CV Diseases	None	None	None	None	None	None
Sanjay Kaul	ACCF/AATS/ SCAI/STS TAVR ECD Writing Committee	Cedars-Sinai Medical Center— Director, Cardiology Fellowship Training Program	None	None	None	None	None	None
Lloyd Klein	Content Reviewer— ACCF Interventional Scientific Council	Rush-Presbyterian-St. Luke's Medical Center— Professor of Medicine	None	None	None	None	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
Joel S. Landzberg	Content Reviewer – ACCF Adult Congenital and Pediatric Cardiology Council	Westwood Cardiology Associates	None	None	None	None	None	None
Ehtisham Mahmud	Content Reviewer— ACCF Interventional Scientific Council	University of California, San Diego— Professor of Medicine and Director,	<ul style="list-style-type: none"> <li>• Phillips Medical</li> </ul>	<ul style="list-style-type: none"> <li>• Medtronic</li> </ul>	None	<ul style="list-style-type: none"> <li>• Abbott Vascular*</li> <li>• Boston Scientific*</li> </ul>	<ul style="list-style-type: none"> <li>• St. Jude Medical</li> </ul>	
James B. McClurken	Content Reviewer— ACCF Surgeon Scientific Council	Temple University Hospital	None	None	None	None	None	None
Geno J. Merli	Content Reviewer— ACCF/AHA/ ACP Task Force on Clinical Competence	Jefferson Center for Vascular Disease— Director; Jefferson Medical	None	None	None	None	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
	and Training	College—  Senior Vice President; Thomas Jefferson University Hospital— Chief Medical Officer						
Issam Moussa	Content Reviewer— ACCF Interventional Scientific Council	Mayo Clinic— Chair, Division of CV Diseases	None	None	None	None	None	None
Brent J. Muhlestein	Content Reviewer— ACCF/AHA/ ACP Task Force on Clinical Competence and Training	Intermountain Medical Center, Cardiology Department— Professor of Medicine	None	None	None	None	None	Defendant, Cardiac Catheterization, 2011
Srihari Naidu	Content Reviewer—	Winthrop University	<ul style="list-style-type: none"> <li>Abbott Vascular</li> </ul>	None	None	None	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
	ACCF Interventional Scientific Council	Hospital— Director, Cardiac Cath Lab						
Hani Najm	Content Reviewer— ACCF Surgeon Scientific Council	National Guard Health Affairs— President, Saudi Heart Association	None	None	None	None	None	None
Jennifer S. Nelson	Official Reviewer— STS	University of Michigan	None	None	None	None	None	None
Ileana Piña	Content Reviewer— ACCF/AHA/ ACP Task Force on Clinical Competence and Training	Montefiore Medical Center— Professor of Medicine and Epidemiology and Population Health	<ul style="list-style-type: none"> <li>• GE HealthCare</li> </ul>	None	None	None	None	None
Stephen Ramee	Official Reviewer— ACC Board of Governors	Ochsner Clinic Foundation— Director, Cardiac	None	None	<ul style="list-style-type: none"> <li>• Boston Scientific*</li> </ul>	<ul style="list-style-type: none"> <li>• Abbott†</li> <li>• Boston Scientific†</li> <li>• Edwards Lifesciences†</li> </ul>	None	None

Peer Reviewer	Representation	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Institutional, Organizational, or Other Financial Benefit	Expert Witness
		Catheterization				<ul style="list-style-type: none"> <li>Medtronic†</li> </ul>		
Richard Smalling	Content Reviewer—ACCF Surgeon Scientific Council	University of Texas Medical School—Jay Brent Sterling Professor of Cardiovascular Medicine	<ul style="list-style-type: none"> <li>AGA Medical</li> </ul>	None	None	<ul style="list-style-type: none"> <li>AGA Medical</li> <li>Cordis</li> <li>E-valve*</li> </ul>	<ul style="list-style-type: none"> <li>AGA Medical*</li> <li>Cordis*</li> </ul>	None
Wilson Y. Szeto	Official Reviewer—STS	University of Pennsylvania	None	None	None	<ul style="list-style-type: none"> <li>Edwards Lifesciences</li> </ul>	None	None
Vinod H. Thourani	Official Reviewer—STS	Emory University	<ul style="list-style-type: none"> <li>Edwards Lifesciences</li> <li>Sorin Medical</li> <li>St. Jude Medical</li> </ul>	None	None	None	None	None

This table represents the relationships of reviewers with industry and other entities that were disclosed at the time of peer review and determined to be relevant. It does not necessarily reflect relationships with industry at the time of publication. A person is deemed to have a significant interest in a business if the interest represents ownership of  $\geq 5\%$  of the voting stock or share of the business entity, or ownership of  $\geq \$10,000$  of the fair market value of the business entity; or if funds received by the person from the business entity exceed 5% of the person's gross income for the previous year. A relationship is considered to be modest if it is less than significant under the preceding definition. Relationships that exist with no financial benefit are also included for the purpose of transparency. Relationships in this table are modest unless otherwise noted. Names are listed in alphabetical order within each category of review.

According to the ACCF, a person has a *relevant* relationship IF: a) The *relationship or interest* relates to the same or similar subject matter, intellectual property or asset, topic, or issue addressed in the *document*; or b) The *company/entity* (with whom the relationship exists) makes a drug, drug class, or device addressed in the *document*, or makes a competing drug or device addressed in the *document*; or c) The *person or a member of the person's household*, has a reasonable potential for financial, professional or other personal gain or loss as a result of the issues/content addressed in the *document*.

\*Significant relationship.

†No financial benefit.

AATS indicates American Association for Thoracic Surgery; ACCF, American College of Cardiology Foundation; SCAI, Society for Cardiovascular Angiography and Interventions; and STS, Society of Thoracic Surgeons.

## REFERENCES

1. ACCF Relationship With Industry and Other Entities: Policies and Procedures for the Development of Guidelines and Other Clinical Documents. [cited 2011; Available from: <http://www.cardiosource.org/Science-And-Quality/Practice-Guidelines-and-Quality-Standards/Relationships-With-Industry-Policy.aspx>.
2. Ruiz, C.E., et al., *Interventional fellowship in structural and congenital heart disease for adults*. JACC Interventions, 2010. **3**(9): p. e1.
3. Feldman, T., C.E. Ruiz, and Z.M. Hijazi, *The SCAI Structural Heart Disease Council: Toward addressing training, credentialing, and guidelines for structural heart disease intervention*. Catheterization and Cardiovascular Interventions, 2010. **76**(4): p. E87-E89.
4. Vassiliades Jr, T.A., et al., *The clinical development of percutaneous heart valve technology: a position statement of the Society of Thoracic Surgeons (STS), the American Association for Thoracic Surgery (AATS), and the Society of Cardiovascular Angiography and Intervention (SCAI)*. Catheterization and Cardiovascular Interventions, 2005. **65**(1): p. 73-79.
5. Herrmann, H.C., et al., *Results of the society of cardiac angiography and interventions survey of physicians and training directors on procedures for structural and valvular heart disease*. Catheterization and Cardiovascular Interventions, 2010. **76**(4): p. E106-E110.
6. Cameron, A., W. Laskey, and W. Sheldon, *SCAI Ad Hoc Task Force on Ethics in Invasive and Interventional Cardiology*. Catheter Cardiovasc Interv, 2004. **61**: p. 157-62.
7. Kouchoukos, N.T., et al., *Guidelines for credentialing of practitioners to perform endovascular stent-grafting of the thoracic aorta*. J Thorac Cardiovasc Surg, 2006. **131**(3): p. 530-2.
8. Beller, G., et al., *28th Bethesda Conference. Task Force 3: Guidelines for credentialing practicing physicians*. Journal of the American College of Cardiology, 1997. **29**(6): p. 1148.
9. *Guidelines for standards in cardiac surgery. Advisory Council for Cardiothoracic Surgery, American College of Surgeons*. Bull Am Coll Surg, 1997. **82**(2): p. 27-9.
10. Vahanian, A., et al., *Transcatheter valve implantation for patients with aortic stenosis: a position statement from the European Association of Cardio-Thoracic Surgery (EACTS) and the European Society of Cardiology (ESC), in collaboration with the European Association of Percutaneous Cardiovascular Interventions (EAPCI)*. Eur J Cardiothorac Surg, 2008. **34**(1): p. 1-8.
11. Peterson, E.D., et al., *Procedural volume as a marker of quality for CABG surgery*. JAMA: the journal of the American Medical Association, 2004. **291**(2): p. 195.
12. Crawford, F.A., et al., *Volume requirements for cardiac surgery credentialing: a critical examination*. The Annals of thoracic surgery, 1996. **61**(1): p. 12-16.
13. Bashankaev, B., S. Baido, and S.D. Wexner, *Review of available methods of simulation training to facilitate surgical education*. Surg Endosc, 2011. **25**(1): p. 28-35.
14. Hislop, S.J., et al., *Simulation case rehearsals for carotid artery stenting*. Eur J Vasc Endovasc Surg, 2009. **38**(6): p. 750-4.
15. Cacho, F., M. Doblare, and G.A. Holzapfel, *A procedure to simulate coronary artery bypass graft surgery*. Med Biol Eng Comput, 2007. **45**(9): p. 819-27.
16. Tedesco, M.M., et al., *Simulation-based endovascular skills assessment: the future of credentialing?* J Vasc Surg, 2008. **47**(5): p. 1008-1; discussion 1014.
17. Seymour, N.E., et al., *Virtual reality training improves operating room performance: results of a randomized, double-blinded study*. Ann Surg, 2002. **236**(4): p. 458-63; discussion 463-4.

18. Bonow, R.O., et al., *2008 Focused Update Incorporated Into the ACC/AHA 2006 Guidelines for the Management of Patients With Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1998 Guidelines for the Management of Patients With Valvular Heart Disease) Endorsed by the Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons*. Journal of the American College of Cardiology, 2008. **52**(13): p. e1.
19. Neily, J., et al., *Association between implementation of a medical team training program and surgical mortality*. JAMA: the journal of the American Medical Association, 2010. **304**(15): p. 1693.
20. Hijazi, Z.M., et al., *Transcatheter Closure of ASDs and PFOs: A Comprehensive Assessment*. 2010: Not Avail.
21. Holmes Jr, D.R., et al., *2012 ACCF/AATS/SCAI/STS Expert Consensus Document on Transcatheter Aortic Valve Replacement*. Journal of the American College of Cardiology, 2012: p. j. jacc. 2012.01. 001v1.
22. Bach, D.S., et al., *Evaluation of patients with severe symptomatic aortic stenosis who do not undergo aortic valve replacement*. Circulation: Cardiovascular Quality and Outcomes, 2009. **2**(6): p. 533-539.
23. Leon, M.B., et al., *Transcatheter Aortic-Valve Implantation for Aortic Stenosis in Patients Who Cannot Undergo Surgery*. New England Journal of Medicine, 2010. **363**(17): p. 1597-1607.
24. Cribier, A., et al., *PERCUTANEOUS TRANSLUMINAL VALVULOPLASTY OF ACQUIRED AORTIC STENOSIS IN ELDERLY PATIENTS: AN ALTERNATIVE TO VALVE REPLACEMENT?* The Lancet, 1986. **327**(8472): p. 63-67.
25. Bashore, T.M. and C.J. Davidson, *Follow-up recatheterization after balloon aortic valvuloplasty*. Journal of the American College of Cardiology, 1991. **17**(5): p. 1188-1195.
26. ANDERSEN, H.R., L.L. KNUDSEN, and J.M. HASENKAM, *Transluminal implantation of artificial heart valves. Description of a new expandable aortic valve and initial results with implantation by catheter technique in closed chest pigs*. European Heart Journal, 1992. **13**(5): p. 704-708.
27. Smith, C.R., et al., *Transcatheter versus Surgical Aortic-Valve Replacement in High-Risk Patients*. New England Journal of Medicine, 2011. **364**(23): p. 2187-2198.
28. Alfieri, O., et al., *The double-orifice technique in mitral valve repair: a simple solution for complex problems*. The Journal of thoracic and cardiovascular surgery, 2001. **122**(4): p. 674.
29. Grube, E., et al., *Percutaneous Aortic Valve Replacement for Severe Aortic Stenosis in High-Risk Patients Using the Second- and Current Third-Generation Self-Expanding CoreValve Prosthesis: Device Success and 30-Day Clinical Outcome*. J Am Coll Cardiol, 2007. **50**(1): p. 69-76.
30. Tamburino, C., et al., *Incidence and Predictors of Early and Late Mortality After Transcatheter Aortic Valve Implantation in 663 Patients With Severe Aortic Stenosis*. Circulation, 2011. **123**(3): p. 299-308.
31. Society of Thoracic Surgery National Database. August 17, 2011]; Available from: <http://www.sts.org/national-database>.
32. American College of Cardiology National Cardiovascular Data Registry. . [cited 2011 August 17]; Available from: <http://www.ncdr.com/WebNCDR/Common/>.
33. Jacobs, J.P., et al., *Successful linking of the Society of Thoracic Surgeons database to social security data to examine survival after cardiac operations*. Ann Thorac Surg, 2011. **92**(1): p. 32-7; discussion 38-9.

34. Topol, E.J., *Failing the public health—rofecoxib, Merck, and the FDA*. New England Journal of Medicine, 2004. **351**(17): p. 1707-1709.
35. Fontanarosa, P.B., D. Rennie, and C.D. DeAngelis, *Postmarketing surveillance—lack of vigilance, lack of trust*. JAMA: the journal of the American Medical Association, 2004. **292**(21): p. 2647.
36. Blackstone, E.H., *Could It Happen Again?* Circulation, 2005. **111**(21): p. 2717-2719.
37. *Interagency Registry for Mechanically Assisted Circulatory Support*. [cited 2011 August 18]; Available from: <http://www.intermacs.org>.